
BAG-1 Antibody

Subcategory: Rabbit Polyclonal Antibody

Cat. No.: 253638

Unit: 0.1 mg

Description:

Bcl-2-associated athanogene 1 (BAG-1) was first identified as an anti-apoptotic bcl-2-binding protein. Later it was found to bind the molecular chaperones Hsp70 and Hsc70 through its carboxy-terminal sequence (termed the Bag domain), resulting in the inhibition of the refolding activity of these chaperones. It is thought that by binding and inhibiting these molecular chaperones, BAG-1 is able to modulate the expression level of proteins requiring chaperones to fold correctly. One such group of proteins that are affected is glucocorticoid receptors. Other reports have suggested that the level of BAG-1 expression correlates with the aggressiveness of various cancers. Multiple isoforms of BAG-1 are known to exist.

Isotype: Rabbit Ig

Applications: E, WB

Species Reactivity: H, M, R

Format: Each vial contains 0.1 ml IgG in PBS pH 7.4 with 0.02% sodium azide. Antibody was purified by immunogen affinity chromatography.

Alternate Names: BAG-1 ; Bcl-2-binding athanogene-1; receptor-associated protein 46; RAP46

Accession No.: NP_004314

Antigen: KLH-conjugated synthetic peptide encompassing a sequence within human BAG-1.

Application Notes: E: 1:500-1:1,000; WB: 1:100-1:500

Storage: Store at -20°C. Minimize freeze-thaw cycles.

Product is guaranteed one year from the date of shipment.

For research use only, not for diagnostic or therapeutic procedures.