

IL-12 Antibody

Subcategory: Rabbit Polyclonal Antibody

Cat. No.: 251278

Unit: 0.1 mg

Description:

Interleukin-12 (IL-12) is a cytokine that can act as a growth factor for activated T and NK cells, enhance the lytic activity of NK/lymphokine-activated Killer cells, and stimulate the production of IFN-gamma by resting PBMC.

Isotype: Rabbit Ig

Applications: E, IHC, WB

Species Reactivity: H, M, R

Format: Each vial contains 0.1 mg IgG in 0.1 ml (1 mg/ml) of PBS pH7.4, 25% glycerol, 5 mg/ml BSA with 0.09% sodium azide. Antibody was purified by immunogen affinity chromatography.

Alternate Names: IL-12; Interleukin-12; Interleukin-12 subunit alpha; IL-12A; IL-12 subunit p35; Cytotoxic lymphocyte maturation factor 35 kDa subunit; CLMF p35; NK cell stimulatory factor chain 1; NKSF1; IL12A; NKSF1; Interleukin-12 subunit beta; IL-12B; IL-12 subunit p40; Cytotoxic lymphocyte maturation factor 40 kDa subunit; CLMF p40; NK cell stimulatory factor chain 2; NKSF2; IL12B; NKSF2

Accession No.: P29459, P29460

Antigen: KLH-conjugated synthetic peptide encompassing a sequence within the center region of human IL-12.

Application Notes: E: 1:500-1:1,000; WB: 1:100-1:500; IHC: 1:100-1:500

Storage: Store at -20°C. Minimize freeze-thaw cycles.

Product is guaranteed one year from the date of shipment.

For research use only, not for diagnostic or therapeutic procedures.